

Abundance *Gammarid amphipods* 1995 – 2014.

Northwest Atlantic United States

March 2016

Prepared for:

Northeast Regional Ocean Council (NROC)

Northeast Ocean Data

northeastoceandata.org

Prepared by:

Marta M Ribera

Eastern Division Conservation Science Office of The Nature Conservancy

99 Bedford Street, 5th Floor

Boston, MA 02111

marta.ribera@tnc.org

1. INTRODUCTION

This data product shows the seasonal abundance per cubic meter (log-transformed) of Gammarid amphipods (a group of zooplankton). This dataset was created by The Nature Conservancy (TNC) using data provided by the Northeast Fisheries Science Center (NEFSC) in Narragansett, RI. This dataset covers offshore waters from North Carolina to northern Maine, and from the coast line until the 1600 meter bathymetry line. The spatial resolution of this layer is 2600 meters.

Source data was collected by the Northeast Fisheries Science Center (NEFSC) as part of their shelf-wide research vessel surveys (Kane 2007, 2011). These surveys are conducted over the continental shelf, from Cape Hatteras (North Carolina) to Cape Sable (Nova Scotia). Plankton samples were collected using a bongo net as part of two types of cruises: broad-scale surveys dedicated to plankton, and trawl and dredge surveys where plankton samples were also collected across the region. Coordinates of sample locations were also collected using a GPS. In the laboratory, zooplankton organisms were sorted, counted, and identified to the lowest possible taxa. NEFSC provided values in abundances by 100m³. All abundance values were divided by 100 to obtain abundance by cubic meter, and values were log-transformed ($\ln(x+1)$) since the distribution of zooplankton values resembled a Poisson distribution (values skewed to smaller abundances). The reason for adding 1 to the abundances prior to calculating the natural log was to prevent irrational values when abundance values are zero.

Sample points were separated in four groups based on the year and season collected: Spring 1995-2004 (1560 points), Fall 1995-2004 (2048 points), Spring 2005-2014 (1338 points), Fall 2005-2014 (1929 points). Seasons were defined to be consistent with meteorological seasons: Spring (March 1st – May 31st); Fall (September 1st – November 30th). Composites of point data samples for each group were interpolated separately using the ArcGIS function “Diffuse interpolation with barriers” to create surfaces. The reason for using this interpolation method was to prevent values being interpolated across land masses (e.g. from North to South of Cape Cod). No data values in this layer represent areas where there were not enough points to interpolate a surface. Finally, layers were clipped to the area of study.

2. PURPOSE

This layer was created to visualize seasonal patterns of zooplankton abundance across space, and to provide the means for comparison between species and between decades. It addresses data gaps and provides the most up-to-date information on zooplankton abundance for the area of study. This supports the Northeast Regional Ocean Council (NROC) efforts to provide datasets to stakeholders and managers to facilitate coastal and ocean planning.

Zooplankton includes a diverse group of free-floating organisms. Zooplankton organisms are secondary consumers, feeding often on phytoplankton organisms. They provide an essential link within the trophic chain, transferring energy from primary producers to predators higher up the trophic chain. *Gammarids* are a group of species of amphipods, which are small shrimp-like organisms. These amphipods are common zooplankton organisms in our area, especially across the water column over Georges Bank. They are part of the diet of juvenile groundfish, including cod and haddock, and many marine mammals.

3. SOURCES AND AUTHORITIES

- Northeast Fisheries Science Center (NEFSC) zooplankton database
- Kane J (2007) Zooplankton abundance trends on Georges Bank, 1977-2004. ICES Journal of Marine Science 64(5):909-91
- Kane J (2011) Inter-decadal variability of zooplankton abundance in the Middle Atlantic Bight. Journal of Northwest Atlantic Fishery Science 43: 81-92

4. DATABASE DESIGN AND CONTENT

Native storage format: ArcGIS File Geodatabase Raster
Columns and Rows: 366, 377

Number of Bands: 1
 Cell Size: 2600 meters (0.03374 degrees)
 Source Type: generic
 Pixel Type: floating point
 Pixel Depth: 32 Bit
 Statistics:

Season	Decade	Minimum	Maximum	Mean	Standard Deviation
SPRING	1995-2004	-5.66325510e-008	3.18210196495056	0.243863272800526	0.450343127713462
	2005-2014	-7.06285945e-008	3.61756896972656	0.23691483477524	0.483050885585818
FALL	1995-2004	3.56008911e-007	5.82439374923706	0.363819098174839	0.629217528693279
	2005-2014	-3.49132811e-008	3.7605741024017	0.27363469071033	0.51108784506464

Dataset Name:
 GammaridsFall1995to2004
 GammaridsFall2005to2014
 GammaridsSpring1995to2004
 GammaridsFall1995to2004

Dataset Status: Complete

5. SPATIAL REPRESENTATION

Reference System: GCS North American 1983
 Horizontal Datum: North American Datum 1983
 Ellipsoid: Geodetic Reference System 1980
 Linear Unit: Meter (1.0)
 Angular Unit: Degree (0.0174532925199433)
 False Easting: 0.0
 False Northing: 0.0
 Central Meridian: 0.0

Geographic extent: -76.45 to -64.101, 33.674 to 46.395

ISO 19115 Topic Category: biology, environment, oceans

Place Names:

Albemarle Sound, Baltimore Canyon, Bay of Fundy, Block Island Delta, Cashes Ledge, Chesapeake Bay, Cholera Bank, Delaware Bay, Georges Bank, Georges Basin, German Bank, Great South Channel, Gulf of Maine, Hudson Canyon, Hydrographer Canyon, Jeffreys Ledge, Jordan Basin, Lake Ontario, Long Island Sound, Mid-Atlantic Bight, Nantucket Shoals, Norfolk Canyon, Northeast Channel, Stellwagen Bank, Southern New England, Wilkinson Basin

Recommended Cartographic Properties:

(Using ArcGIS ArcMap nomenclature)

Stretch, Standard Deviations (2.5), Prediction color ramp.

To compare different Gammarid layers using the same scale, use statistics from Custom settings (reference values from Fall 1995-2004):

Min: 3.5600891123976908e-007

Max: 5.8243937492370605

Mean: 0.36381909817483987

Std Dev.: 0.62921752869327907

Scale range for optimal visualization: 1:5,000,000

6. DATA PROCESSING

Processing environment: Microsoft Windows 7 Professional, Service Pack 1; ESRI ArcGIS 10.2.2, extensions: Geostatistical Analyst, Spatial Analyst; Microsoft Office 2010

	Process Steps Description
1	In Excel, database was queried for selected decade, season, and species. Abundances were divided by 100 to obtain abundances by 1m ³ . Finally, results were log-transformed (ln+1).
2	Plotted results in ArcGIS. Point data was interpolated into raster surface using DIFFUSION INTERPOLATION WITH BARRIERS from the Geostatistical Analyst extension. Used polygon shapefile of US states as absolute barriers, and bandwidth of 20,000m.
3	Layer was clipped to area of study using EXTRACT BY MASK function from the Spatial Analyst extension.

7. QUALITY PROCESS

Attribute Accuracy: The accuracy of the data is a result of the accuracy of the source data provided by NEFSC.

Logical Consistency: These data are believed to be logically consistent.

Completeness: This layer is a composite from all available data samples within each season and decade. Areas with data gaps signify locations without enough samples to interpolate a surface.

Positional Accuracy: The accuracy of the data is a result of the accuracy of the GPS units used when samples were collected. Also, interpolations of point data samples to surfaces may create inaccuracies in certain areas.

Timeliness: Based on data from January 1995 - December 2014.

Use restrictions: The Nature Conservancy (TNC) compiled this data set from publicly available data sources and this data is freely distributable without permission from this TNC resource office. This data set must be cited on all electronic and hard copy products using the language of the Data Set Credit. Use and analysis of the geographic data are limited by the scale at which the data was collected and mapped, and that, as a regional analysis, it is not intended for site level decisions. The Nature Conservancy shall not be held liable for improper or incorrect use of the data described and/or contained herein. Any sale, distribution, loan, or offering for use of these digital data, in whole or in part, is prohibited without the approval of the Nature Conservancy. The use of these data to produce other GIS products and services with the intent to sell for a profit is prohibited without the written consent of the Nature Conservancy. All parties receiving these data must be informed of these restrictions. The Nature Conservancy shall be acknowledged as data contributors to any reports or other products derived from these data.

Access Constraints: The Nature Conservancy reserves all rights in data provided. All data are provided as is. This is not a survey quality dataset. The Nature Conservancy makes no warranty as to the currency, completeness, accuracy or utility of any specific data. This disclaimer applies both to individual use of the data and aggregate use with other data. It is strongly recommended that careful attention be paid to the contents of the metadata file associated with these data.

Distribution Liability: Neither The Nature Conservancy nor any of the ecoregional planning team participants makes any warranty, expressed or implied as to the use or appropriateness of use of the enclosed data, nor are there warranties of merchantability or fitness for a particular purpose or use. No representation is made as to the currency, accuracy or completeness of the information in this dataset or of the data sources on which it is based. Neither the Nature Conservancy nor any of the ecoregional planning team participants shall be liable for any lost profits or consequential damages, or claims against the user by third parties.